#### Section 7.1: RATIONAL EXPRESSIONS AND THEIR SIMPLIFICATION

When you are done with your homework you should be able to...

- $\pi$  Find numbers for which a rational expression is undefined
- $\pi$  Simplify rational expressions
- $\pi$  Solve applied problems involving rational expressions

WARM-UP:

1. Factor:

$$x^3 - 8x^2 + 2x - 16$$

2. Solve:

$$2x^2 - x - 10 = 0$$

## EXCLUDING NUMBERS FROM RATIONAL EXPRESSIONS

| A                             | expression is the | of two                     |  |  |  |  |
|-------------------------------|-------------------|----------------------------|--|--|--|--|
| Rational expressions indicate |                   |                            |  |  |  |  |
| and division by               | is                | This means that we         |  |  |  |  |
|                               |                   | any value or values of the |  |  |  |  |
| that make a                   |                   |                            |  |  |  |  |

Example 1: Find all numbers for which the rational expression is undefined:

a. 
$$\frac{5}{x}$$

b. 
$$\frac{x+1}{x-4}$$

c. 
$$\frac{8x-40}{x^2+3x-28}$$

d. 
$$\frac{x-12}{x^2+4}$$

## SIMPLIFYING RATIONAL EXPRESSIONS

| A |            | is |     | if its |
|---|------------|----|-----|--------|
|   | and        | ho | ave | common |
|   | other than | or |     |        |

## FUNDAMENTAL PRINCIPLE OF RATIONAL EXPRESSIONS

If \_\_\_\_\_, and \_\_\_\_ are \_\_\_\_ and \_\_\_\_ and \_\_\_\_ and \_\_\_\_ are \_\_\_\_

# STEPS FOR SIMPLIFYING RATIONAL EXPRESSIONS

1. \_\_\_\_\_ the \_\_\_\_ and the \_\_\_\_

completely.

2. \_\_\_\_\_ both the \_\_\_\_ and the

\_\_\_\_\_ by any \_\_\_\_\_\_.

3. \_\_\_\_\_ the \_\_\_\_\_ in the \_\_\_\_\_ equation.

Example 2: Simplify:

a. 
$$\frac{4x-64}{16x}$$

b. 
$$\frac{6y+18}{11y+33}$$

c. 
$$\frac{x^2 - 12x + 36}{4x - 24}$$

d. 
$$\frac{x^3 + 4x^2 - 3x - 12}{x + 4}$$

$$e. \ \frac{x+5}{x-5}$$

f. 
$$\frac{x^3-1}{x^2-1}$$

# SIMPLIFYING RATIONAL EXPRESIONS WITH OPPOSITE FACTORS IN THE NUMERATOR AND DENOMINATOR

The \_\_\_\_\_ of two \_\_\_\_ that have \_\_\_\_ is \_\_\_\_ is \_\_\_\_ .

Example 3: Simplify:

a. 
$$\frac{x-3}{3-x}$$

b. 
$$\frac{9x-15}{5-3x}$$

c. 
$$\frac{x^2-4}{2-x}$$

#### **APPLICATION**

A company that manufactures small canoes has costs given by the equation

$$C = \frac{20x + 20000}{x}$$

in which x is the number of canoes manufactured and  $\mathcal C$  is the cost to manufacture each canoe.

a. Find the cost per canoe when manufacturing 100 canoes.

b. Find the cost per canoe when manufacturing 10000 canoes.

c. Does the cost per canoe increase or decrease as more canoes are manufactured?