

When you finish your homework you should be able to...

- π Use Sigma Notation to Write and evaluate a Sum
- π Understand the Concept of Area
- π Approximate the Area of a Plane Region
- π Find the Area of a Plane Region Using Limits

Warm-up: Evaluate the following limits.

1. $\lim_{x \rightarrow \infty} \frac{25x^2 - 5x + 3}{x^2}$

2. $\lim_{x \rightarrow \infty} \left(\frac{2x^3 + 6x^2}{3x^3} + 10 \right)$

SIGMA NOTATION

The sum of n terms a_1, \dots, a_n is written as $\sum_{i=1}^n a_i = a_1 + a_2 + a_3 + \dots + a_n$ where i is the **index of summation**, a_i is the **i th term** of the sum, and the **upper and lower bounds of summation** are n and 1.

Example 1: Find the sum.

a. $\sum_{i=1}^5 i$

b. $\sum_{i=1}^4 i^2$

c. $\sum_{i=1}^3 \sqrt{i}$

Summation Properties

1.
$$\sum_{i=1}^n ka_i = k \sum_{i=1}^n a_i$$

2.
$$\sum_{i=1}^n (a_i \pm b_i) = \sum_{i=1}^n a_i \pm \sum_{i=1}^n b_i$$

Example 2: Find the sum.

a.
$$\sum_{i=1}^4 2$$

b.
$$\sum_{i=1}^3 (1 - i^2)$$

Theorem: Summation Formulas

1.
$$\sum_{i=1}^n c = cn$$

2.
$$\sum_{i=1}^n i = \frac{n(n+1)}{2}$$

3.
$$\sum_{i=1}^n i^2 = \frac{n(n+1)(2n+1)}{6}$$

4.
$$\sum_{i=1}^n i^3 = \frac{n^2(n+1)^2}{4}$$


Example 3: Evaluate the following sums.

a.
$$\sum_{i=1}^n (3i - i^2)$$


b.
$$\sum_{i=1}^n (6i + 4i^3)$$

Area


In Euclidean geometry, the simplest type of plane region is a _____. The definition for the area of a rectangle is _____. From this definition, you can develop formulas for the areas of many other plane regions such as triangles. To determine the area of a triangle, you can find a _____ whose area is _____ that of the _____. Once you know how to find the area of a triangle, you can determine the area of any _____ by subdividing the polygon into _____ regions.



Parallelogram




Hexagon



Polygon

Figure 5.6


Finding the areas of regions other than polygons is more difficult. The ancient Greeks were able to determine formulas for the areas of some general regions by the _____ method. Essentially, the method is a _____ process in which the area is _____ between two polygons—one _____ in the region and one _____ about the region.




ARCHIMEDES (287–212 B.C.)

Archimedes used the method of exhaustion to derive formulas for the areas of ellipses, parabolic segments, and sectors of a spiral. He is considered to have been the greatest applied mathematician of antiquity.

See LarsonCalculus.com to read more of this biography.



$n = 6$



$n = 12$

The exhaustion method for finding the area of a circular region

Figure 5.7

$$s(n) \leq (\text{Area of region}) \leq S(n)$$




Figure 5.11

Theorem: Limits of the Lower and Upper Sums

Let f be continuous and nonnegative on the interval $[a, b]$. The limits as $n \rightarrow \infty$ of both the lower and upper sums exist and are equal to each other. That is,


$$\begin{aligned} \lim_{n \rightarrow \infty} s(n) &= \lim_{n \rightarrow \infty} \sum_{i=1}^n f(m_i) \Delta x \\ &= \lim_{n \rightarrow \infty} \sum_{i=1}^n f(M_i) \Delta x \\ &= \lim_{n \rightarrow \infty} S(n) \end{aligned}$$

where $\Delta x = \frac{b-a}{n}$ and $f(m_i)$ and $f(M_i)$ are the minimum and maximum values of f on the subinterval.

Example 4:


a. Find the upper sum from

$x = 2$ to $x = 6$.



b. Find the lower sum from

$x = 2$ to $x = 6$.



Definition of an Area in the Plane

Let f be continuous and nonnegative on the interval $[a, b]$. The area of the region bounded by the graph of f , the x -axis, and the vertical lines $x = a$ and $x = b$ is

$$\text{Area} = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(c_i) \Delta x, \quad x_{i-1} \leq c_i \leq x_i \quad \text{where } \Delta x = \frac{b-a}{n},$$

right endpoint: $c_i = a + i\Delta x$, left endpoint: $c_i = a + (i-1)\Delta x$

Example 5: Find the area of the region bounded by the graph $f(x) = x^3$, the x -axis, and the vertical lines $x = 0$ and $x = 1$.