

When you are done with your homework you should be able to...

- $\pi$  Understand and use Rolle's Theorem
- $\pi$  Understand and use the Mean Value Theorem

**Warm-up:** Locate the global extrema of the function  $f(x) = x^3 - 12x$  on the closed interval  $[0, 4]$ .

### ROLLE'S THEOREM


The \_\_\_\_\_ Value Theorem states that a continuous function on a closed interval \_\_\_\_\_ must have both a \_\_\_\_\_ and a \_\_\_\_\_. Both of these values, however, can occur at the \_\_\_\_\_. Rolle's Theorem gives conditions that guarantee the existence of an extreme value in the \_\_\_\_\_ of a closed interval.

### THEOREM: ROLLE'S THEOREM

Let  $f$  be continuous on the closed interval  $[a, b]$  and differentiable on the open interval  $(a, b)$ . If  $f(a) = f(b)$  then there is at least one number  $c$  in  $(a, b)$  such that  $f'(c) = 0$ .

Example 1: Determine whether Rolle's Theorem can be applied to  $f$  on the closed interval  $[a, b]$ . If Rolle's Theorem can be applied, find all values of  $c$  in the open interval  $(a, b)$  such that  $f'(c) = 0$ .

a.  $f(x) = \cos 2x, [-\pi, \pi]$



b.  $f(x) = x^2 - 5x + 4, [1, 4]$

